



# YTOLOGY

Premed 2018 - JU

Sheet

Slides

Number

2

Done by:

Laila Nazzal

Corrected by:

Obada froukh

Doctor

Diala abu Hassan

In the previous lecture we have considered the non-covalent interactions that are classified into:

van der  
waals

ionic  
interactions

hydrophobic  
interactions

hydrogen  
interactions

### Hydrogen interactions

- Hydrogen interactions are a special type of **ionic interactions**
- They are of the strongest non-covalent interactions

**Remember:** (hydrogen is not a highly electronegative atom i.e. it does not have a high ability to attract the electrons. Therefore, mostly it would have a partial positive charge when it is bonded to a high electronegative atom such like oxygen, nitrogen, etc).

#### So, how does the hydrogen bond form ?

As the name implies, hydrogen is involved for sure

NOW, if hydrogen is bonded to such high electronegative atoms in a molecule, it gets a partial positive charge and if there is a highly electronegative atom in ANOTHER molecule that's charged with a partial negative charge, then the partial negative charge (in the 2nd molecule is attracted to the partial positive charge on hydrogen in the 1st molecule) resulting in the hydrogen bond

-- → And that's why it's considered a special type of ionic interactions (attraction of charges)

The hydrogen that contributes to the formation of hydrogen bond = hydrogen bond **DONOR**.

The other atom = hydrogen bond **ACCEPTOR**.


So, to form a hydrogen bond you need a donor and an acceptor

i.e. two accepters  
without a donor  
cannot form a  
hydrogen bond.


→ as shown in the  
figure:

In the water  
molecule,

- There are two donors – two hydrogen atoms connected to a highly electronegative atom-
- One acceptor-oxygen- (so it has a high ability to form a large number hydrogen bonds)
- The length of the hydrogen bond is 0.177 nm .
- The length of the covalent bond(OH) is 0.0965 nm .  
Which is shorter?  
-covalent .  
So, which is stronger?  
-covalent .



### Hydrogen bond



Lehninger 5<sup>th</sup> edition  
Dr. DIALA Abu-Hassan




FIG. 4.3. Hydrogen bonds between water molecules. The oxygen atoms are shown in black.

Mark's 2012

12




**/covalent bonds are stronger than non-covalent interactions generally/**

Generally, What's the importance of non-covalent interactions ,particularly hydrogen bonds?

1. They hold the structure of very large molecules such as DNA and Proteins –they contribute to the final 3D shape of a protein-

### Noncovalent Interactions



### Examples:

- **collagen**

actually, it's the most abundant protein in the body, and is found in hair and skin. Also, it has more than one type as it is found in bones and many other tissues.

Its fibril form is composed of 3 helices winded around each other

-What holds these together? **HYDROGEN BONDS**.

As a protein it has a structural role so its main function is to give **support** in bone..skin ,etc.

- **DNA molecule:**


The Double helix : two strands are held together by the base pairing between (C and G), (A and T), the base pairing is hydrogen bonds!

i.e.the whole strands are held together by the hydrogen bonds which leads to the final structural and functional shape of DNA .

- **Water**

The hydrogen bond contributes to the Physical properties of water, such as:

- ❖ Adhesion
- ❖ Melting and boiling point :if you compare the structure of a water molecule to other molecules, relatively its high .
- ❖ Cohesion/Surface tension
- ❖ Expansion when frozen
- ❖ **Its specific heat (or heat capacity) is higher** than other liquids .



Remember: specific heat is the amount of energy needed to increase the temperature of 1 gram of water molecules(or substance) about 1 degree



-and this is important, as you move to a cold place for example, you don't have a large change in the temperature of your body due to the high specific heat of water inside it

❖ It acts as a solvent for many solutes .

**Solving:** physical process in which molecules of solute enter between the inter molecular spaces of the water molecules and bind with them via the hydrogen bonds.

Eg: solving salt (NACL) in water



As shown in the figure aside:

Water molecules start to interfere ions (Na+ and Cl-) until they reach to the point that each ion is surrounded

with water molecules forming **THE HYDRATION SHELL** which is composed of the ion(in the middle) surrounded by water molecules .

- NOTE that the water molecules direct themselves in a clever way, how?

### Water as a solvent



Dr. Diala Abu-Hassan

Lehninger 5<sup>th</sup> edition

Mark's 2012

14

Look at Na+, water directs its oxygen atom(which has a partial negative charge) towards it, whereas it directs the positive charge of the hydrogen towards the negative charge of Cl-.


Notes:

- **Not** only ionic compounds are dissolved in water, for example RCOR( ketone) as shown in the fig aside .

- Is it a hydrogen bond donor, acceptor or both?

It's an **acceptor**.

- Does It form a hydrogen bond with molecules of the same group (ketones)??



H-bonds between water and polar molecules. *R* denotes additional atoms.

No, as said, an acceptor with an acceptor does not form a hydrogen bond.

**That's why the boiling and melting point of acetone –in this example-is very low!**

- Just to keep in mind how its boiling point is very low. Remember for example that Acetone is used in removing the nail paint and it volatilize very fast ^^

-For solvation to be formed it needs to form a hydrogen bonds with water, where hydrogens of water as donor and the molecule (ketone) as an acceptor.

-----

Well, does **any** hydrogen atom form a hydrogen bond, what do you think?

Actually not all hydrogen atoms form hydrogen bonds. For example, hydrogen attached to carbon, doesn't have a partial positive charge so it can't form a hydrogen bond 😊

-----

## Properties of Noncovalent Interactions

1. Reversible
2. Relatively weak. 1-30 kj/mole vs. 350 kj/mole in C—C bond
3. Molecules interact and bind specifically.

**What are the general properties of all non-covalent interactions ?**

# Noncovalent Interactions


-Noncovalent forces significantly contribute to the structure, stability, and functional competence of macromolecules in living cells.

- Can be either attractive or repulsive.

-Involve interactions both within the biomolecule and between it and the water of the surrounding environment.

-Involve interactions both within the biomolecule and between it and the water of the surrounding environment.

Let's discuss more:



1) **Reversible** : they form and break all the time because molecules are moving.

(Remember, the main factors that affect the formation of non covalent interactions are **DISTANCE AND ORIENTATION** .

i.e. if they move away from each other or are in an improper orientation , bond is broken .

2) They are **relatively weak**: specifically if we compare them to covalent bonds .

3) They interact and bind **specifically**: we have certain conditions, +Ve charge with -Ve charge are attracted, -ve and -ve repulse each other, hydrogen bond has specific conditions to form and so on .

- **Non-covalent interactions rely on the number of bonds rather than the strength of each single bond**, for them to contribute to the structure of DNA and proteins, we need a large number of bonds .

# Covalent vs Noncovalent bonds

	Type of Bond	Energy	
		(kJ mol <sup>-1</sup> )	(kcal mol <sup>-1</sup> )
Covalent Bonds (Strong)	O—H	460	110
	H—H	416	100
	C—H	413	105
Noncovalent Bonds (Weaker)	Hydrogen bond	20	5
	Ion-dipole interaction	20	5
	Hydrophobic interaction	4–12	1–3
	Van der Waals interactions	4	1

\*Note that two units of energy are used throughout this text. The kilocalorie (kcal) is a commonly used unit in the biochemical literature. The kilojoule (kJ) is an SI unit and will come into wider use as time goes on. The kcal is the same as the "Calorie" reported on food labels.

Dr. Diala Abu-Hassan

Campbell 6th edition



If you compare -in terms of strength-  
covalent and non-covalent bonds you will  
see that covalent bonds are stronger for  
sure ^^

# Carbohydrates

What are carbohydrates ? ❖

**Poly hydroxy aldehydes or ketones naturally occurring compounds -organic molecule.**

- *Poly*: many .

-*Hydroxy* : contains hydroxyl groups .

-*Aldehydes* : contains carbonyl group (CO) at carbon number one .

-*Ketones* : contains carbonyl group (CO) at carbon- at least- number two .

-*ose* : means this is a sugar .

-*Ketose* : ketone sugar .


-*Aldose* : aldehyde sugar .

**\* How many carbon are present in the smallest aldehyde sugar or ketone sugar?**

**Three**


**Ok, why**

Let's think about an aldehyde with two carbons, one of them will have the carbonyl group and the 2<sup>nd</sup> will have the hydroxyl group .



The simplest  
ketose sugar is :  
dihydroxyacetone




But  
remember!  
Sugars must  
contain a  
carbonyl group  
and MORE  
than one  
hydroxyl group  
in its structure.



The simplest  
aldose sugar is :  
glyceraldehyde

# Carbohydrates

- ✓ Include both simple monosaccharides and the complex polysaccharides
- ✓ General formula for simple sugars  $(CH_2O)_n$ ,  $n=3-7$  mostly
- ✓ 3, 5, 6 are the most common
- ✓ Cyclization
- ✓  $\alpha$  and  $\beta$  forms on carbon 1



as shown in the figure above:

**Carbohydrates are classified according to their structure:**

- 1) linear form .
- 2) cyclized form (Rings) .

Note: Ribose sugar contains 5 carbons (pentose)," pent" refers to 5 and "ose" to sugar, it is an aldose because of the presence of carbonyl group on carbon number 1

Mostly this sugar (pentose) interact to form rings( **carbon number 1 with hydroxyl group of carbon 4 -before the last carbon- in aldehydes** ), if it is a ketose then carbon 2 with 4

note: Linear form is occasionally present


the OH groups which are in the linear form on the right side of carbons they will be downwards whereas the OH groups which are on the left side as in glucose on 3C will be upwards

Right → down the ring

Left → above the ring

Keep in mind that alpha or beta form depends on the anomeric carbon- **for example C number 1 in aldehydes-**

**The smallest sugar that can form a ring has 4 carbons .**



Carbohydrates - Cyclization o... dept.harpercollege.edu

**Where does Hydrogen of the OH group on the anomeric carbon come from ?**

--> because of this interaction we need to break one of its bonds (double bonds) on **CO**, for the oxygen it needs another bond to be saturated so that it binds to a Hydrogen and this creates a hydroxyl group on carbon number one.

*\*Will this OH group be upward or downward ?*

it has the two orientations, it can be either upward ( $\beta$ ) or downward ( $\alpha$ ).

- each sugar molecule is present in all forms alpha, beta and linear and there is equilibrium between those forms and it converts from one form to another.

Note: moving from ( $\alpha$ ) to ( $\beta$ ) or vice versa we need to pass through the linear form.

*look at the structures again, suppose that I give you 100% time and we want to divide it between alpha, beta and linear, how will you divide it ?*

Actually linear is the least stable due to the functional groups it has

BETA is about 64%

ALPHA is about 36%

Why??

Beta is more stable, as it has its OH group directed UPWARDS, and the one next to it is directed downwards, so I REDUCED the REPULSION between them (much more than alpha)

**\*most of the time beta sugars are more stable BUT you should pay attention to carbon number 2 or 3 – depending on its type.. aldehyde or ketone- and see the direction of the OH group on it (and sometimes other), in the case of aldehydes: if OH is downwards then beta sugars are more stable, if its upwards then alpha sugars are more stable\*.**

**More explanation: we must keep larger molecules further away from each other (eg: keep away OH FROM CH2OH in ketoses and OH FROM OH in aldehydes)**


DNA molecule has one form of sugar –which is beta deoxyribose-

Here, it makes a difference  
Whether you choose alpha or beta  
Because DNA is a very large molecule  
And the stereochemistry plays a role here.

On the other hand, if I have a glucose molecule and I want to metabolize it (extract energy), it doesn't matter whether its alpha or beta.

## Disaccharides

- ✓ **Connecting monosaccharides:**
  - The OH group on anomeric carbon of the 1<sup>st</sup> molecule is going to interact With OH group of the other carbon (usually not the anomeric one) .



# Interaction happens between first molecule as a hemiacetal .

Hemiacetal: it's a molecule that has an (OH) group and (OR) group on the same carbon [look at the figure aside before the reaction occurs, the anomeric carbon - 1<sup>st</sup> carbon]

- Interacting to alcohol (OH group of carbon 4 in the 2<sup>nd</sup> molecule)
- Hemiacetal + alcohol = acetal

Heme  
hemiacetal  
has one  
and after  
is finished it  
as it has 2  
☺


- ✓ Dehydration reaction
- ✓ 2 residues=disaccharide
- ✓ A few residues=oligosaccharide
- ✓ Hundreds or thousands residues=polysaccharide

means half,  
because it  
(OR) group  
the reaction  
is an acetal  
(OR) groups

## Polysaccharides

✓ Storage polysaccharides are starch (plants) and glycogen (animal)

✓ Cellulose is a structural sugar



\*Just in case you get confused about alpha and beta and which one is digested, think about it in this way, as I mentioned before if you want to metabolize glucose for example in order to get energy, it makes no difference whether its alpha or beta, the point is that the ENZYME which recognize  $\alpha(1-4)$  glycosidic linkage does NOT recognize beta ones, and that's why cellulose is not digested in our bodies for example (due to the absence of the enzyme cellulase) .


-Notice the shape of linkage in alpha(it looks like 7) but in beta it looks like (N letter) .

### **Structural polysaccharides**

### **Cellulose:**

- Its made up of  $\beta$  glucose and connected with  $\beta(1-4)$  glycosidic linkage .
- Cellulose fibers are found in fruits, vegetables and trees .
- It is a structural polysaccharide .
- Linear, does not have branches (which makes the whole surface area open for hydrogen bonding  $\rightarrow$  mechanical strength) .

## Cellulos




- Animals lack cellulases enzymes that
- Cellulases are found in the bacteria t of insects and grazing animals, such :


## Starch

Types of starches

Amylose



Amylopectin



Dr. Dalia Abu-Hassan

22

## Storage polysaccharides

### 1) Starch :

a) Amylopectin

- It's a component of starch(storage polysaccharide in plants).
- It can be digested as it has alpha one four linkage .
- Branched .
- Major.

b) Amylose


- unbranched
- 20% of starch (minor) .
- Linear molecule that forms a helix .

### 2) Glycogen

- Glucose as a monomer with  $\alpha(1-4)$  glycosidic linkage .
- Much more branching with a layers of it .

- Storage polysaccharide in human body and animal cells .
- Stored in liver and muscles .

### Amylopectin versus glycogen



Amylopectin


Glycogen

- Branching
- Source


### *What is the importance(roles) of carbohydrates?*

## Glycogen

- Glycogen is more branched than starch
- Branch points occur about every 10 residues in glycogen
- $\alpha$  (1-4) linkage in main chain
- $\alpha$  (1-6) linkage at branches
- The average chain length is 13 glucose residues
- 12 layers of branching



Glycogen



23

Dr. Diala Abu-Hassan

glyx  
protein  
cytoskeleton

# Roles of Carbohydrates

## Major energy sources

Starting material for the **synthesis of other cell constituents**

Major **nutrients** of the cells

Oligosaccharides play a key role in **cell–cell interactions, adhesion, immune recognition, protein targeting and protein folding.**

Polysaccharides are essential **structural components** of several classes of organisms such as cellulose (a major component of grass and trees)

Dr. Diala Abu-Hassan

25

Let's end this sheet with some examples☺ :

**1) structural proteins : for example, Glycocalyx around membranes that have a structural role acting as a layer around the whole cell that protect .**

### **2) cell-cell recognition :**

The difference between blood groups in humans body **is** the sugars that come out of the red blood cells.

### **3) Immune recognition**

About immune recognition: when we get infection by bacteria or virus, the immune system recognize it as an antigen through sugars that are on cells surface.

### **4) protein targeting:**

After synthesizing proteins and modifying them, they are packaged and sent to its final destination, so how does it know that it should go to plasma membrane and not mitochondria for example? By **labeling**, and one type of labeling (tagging) is by using sugars .

### **5) protein folding:**

Forming a 3D shape is contributed by sugars if it is a **glycoprotein**.

- The first thing that your body start to digest to get the energy is sugar .
- The glycogen storage in your body is enough to supply you with energy for about 12 hours if you are fasting and then we start to take our energy from other sources .
- Why is it recommended to eat fibers( cellulose) if you want to lose weight or you have constipation?  
It activate the muscles and increases the contraction .
- Furthermore, it acts as a meshwork that catches toxins, cancerogenic materials, extra cholesterol and others.

(لَوْ أَنَّ النَّاسَ كُلَّ مَا اسْتَصْعَبُوا أَمْرًا تَرَكَوهُ، مَا قَامَ لِلنَّاسِ دُنْيَا وَلَا دِين)

عُمر بن عبد العزيز-